arXiv:0811.2487v1 [math-ph] 16 Nov 2008

The Number of Supertraces on the Superalgebra of Observables of Rational Calogero Model based on the Root System

S.E. Konstein*[†]

I.E.Tamm Department of Theoretical Physics,
P. N. Lebedev Physical Institute,
119991, Leninsky Prospect 53, Moscow, Russia.

and

R. Stekolshchik[‡]

ECI Telecomm Ltd.,
Osivim 30, Petah-Tikva, Israel.

Abstract

In the Coxeter group $W(R)$ generated by the root system R, let $Q(R)$ be the number of conjugacy classes having no eigenvalue -1. The superalgebra $H_{W(R)}$ of observables of the rational Calogero model based on the root system R possesses $Q(R)$ supertraces. The numbers $Q(R)$ are determined for all irreducible root systems (hence for all root systems).

---

*E-mail: konstein@lpi.ru

[†] This work was supported by the RFBR (grant No. 08-02-01118 (S.K.)), and by the grant LSS-1615.2008.2 (S.K.).

[‡] E-mail: rs2@biu.013.net.il

1 The superalgebra of observables

The superalgebra $H_{W(R)(\nu)}$ of observables of the rational Calogero model based on the root system R is defined in the following way.

For any nonzero $\vec{v} \in V = \mathbb{R}^N$ define the reflections $R_{\vec{v}}$ as follows:

$$R_{\vec{v}}(\vec{x}) = \vec{x} - 2\frac{(\vec{x},\, \vec{v})}{(\vec{v},\, \vec{v})}\vec{v} \qquad \text{for any } \vec{x} \in V. \tag{1}$$

Here $(\cdot, \cdot)$ stands for the inner product in V, i.e., $(\vec{x},\, \vec{y}) = \sum_{i=1}^N x_i y_i$, where the x_i are the coordinates of vector $\vec{x}$: $x_i \overset{def}{=} (\vec{x},\, \vec{e}_i)$, and the vectors $\vec{e}_i$ constitute an orthonormal basis in V: $(\vec{e}_i,\, \vec{e}_j) = \delta_{ij}$. The reflections (1) have the following properties

$$R_{\vec{v}}(\vec{v}) = -\vec{v}, \qquad R_{\vec{v}}^2 = 1, \qquad (R_{\vec{v}}(\vec{x}),\, \vec{u}) = (\vec{x},\, R_{\vec{v}}(\vec{u})), \quad \text{for any } \vec{v},\, \vec{x},\, \vec{u} \in V.$$

Definition 1. *The finite set of vectors $R \subset V$ is a root system if R is $R_{\vec{v}}$-invariant for any $\vec{v} \in R$ and the group $W(R)$ generated by all reflections $R_{\vec{v}}$ with $\vec{v} \in R$ (called a Coxeter group) is finite.*

In this paper, we consider noncrystallographic irreducible root systems (H_3, H_4 and $I_2(n)$) together with crystallographic ones (A_n, B_n, C_n, D_n, E_n, F_4, G_2) [1, 2].

Let $\mathcal{H}^\alpha$ ($\alpha = 0, 1$) be two copies of V with orthonormal bases a_i^α ($i = 1, \ldots, N$), respectively. For every vector $\vec{v} = \sum_{i=1}^N v_i \vec{e}_i \in V$ let $v^\alpha \in \mathcal{H}^\alpha$ be the vectors $v^\alpha = \sum_{i=1}^N v_i a_i^\alpha$, so the bilinear forms on $\mathcal{H}^\alpha \otimes \mathcal{H}^\beta$ can be defined as $(x^\alpha,\, y^\beta) = (\vec{x},\, \vec{y})$, where $\vec{x},\, \vec{y} \in V$ and $x^\alpha,\, y^\alpha \in \mathcal{H}^\alpha$ are their copies. The reflections $R_{\vec{v}}$ act on $\mathcal{H}^\alpha$ as follows

$$R_{\vec{v}}(h^\alpha) = h^\alpha - 2\frac{(h^\alpha,\, v^\alpha)}{(\vec{v},\, \vec{v})}v^\alpha, \qquad \text{for any } h^\alpha \in \mathcal{H}^\alpha.$$

Thus, the $W(R)$-action on the spaces $\mathcal{H}^\alpha$ is defined.

Let ν be a set of constants $\nu_{\vec{v}}$ with $\vec{v} \in R$ such that $\nu_{\vec{v}} = \nu_{\vec{w}}$ if $R_{\vec{v}}$ and $R_{\vec{w}}$ belong to one conjugacy class of $W(R)$.

Definition 2. *$H_{W(R)}(\nu)$ is the algebra of polynomials in the a_i^α with coefficients in the group algebra $\mathbb{C}[W(R)]$ subject to the relations*

$$g h^\alpha = g(h^\alpha) g \quad \textit{for any } g \in W(R), \quad \textit{and } h^\alpha \in \mathcal{H}^\alpha$$

$$[h_1^\alpha, h_2^\beta] = \varepsilon^{\alpha\beta}\left((\vec{h}_1,\, \vec{h}_2) + \sum_{\vec{v}\in R} \nu_{\vec{v}} \frac{(\vec{h}_1,\, \vec{v})(\vec{h}_2,\, \vec{v})}{(\vec{v},\, \vec{v})} R_{\vec{v}}\right) \textit{ for any } h_1^\alpha,\, h_2^\alpha \in \mathcal{H}^\alpha, \tag{2}$$

where $\varepsilon^{\alpha\beta}$ is the antisymmetric tensor, $\varepsilon^{01} = 1$.

This algebra is associative because it has faithful representation via Dunkl differential-difference operators [3] acting on the space of infinitely smooth functions on V. Namely, let

$$D_i = \frac{\partial}{\partial x_i} + \frac{1}{2}\sum_{\vec{v}\in R} \nu_{\vec{v}} \frac{v_i}{(\vec{x},\, \vec{v})}(1 - R_{\vec{v}})$$

and [4, 5]

$$a_i^\alpha = \frac{1}{\sqrt{2}}(x_i + (-1)^\alpha D_i), \quad \alpha = 0, 1. \tag{3}$$

The reflections $R_{\vec{v}}$ transform the deformed creation and annihilation operators (3) as vectors:

$$R_{\vec{v}} a_i^\alpha = \sum_{j=1}^{N} \left(\delta_{ij} - 2\frac{v_i v_j}{(\vec{v},\, \vec{v})} \right) a_j^\alpha R_{\vec{v}}.$$

Since $[D_i,\, D_j] = 0$ [3], it follows that

$$[a_i^\alpha, a_j^\beta] = \varepsilon^{\alpha\beta} \left(\delta_{ij} + \sum_{\vec{v}\in R} \nu_{\vec{v}} \frac{v_i v_j}{(\vec{v},\, \vec{v})} R_{\vec{v}} \right),$$

which manifestly coincides with (2).

The commutation relations (2) suggest to define the *parity* π by setting:

$$\pi(a_i^\alpha) = 1 \text{ for any } \alpha,\, i, \qquad \pi(g) = 0 \text{ for any } g \in W(R)$$

and consider $H_{W(R)}(\nu)$ as a superalgebra.

We call $H_{W(R)}(\nu)$ *the superalgebra of observables of the Calogero model based on the root system* R.

Obviously, $\mathbb{C}[W(R)]$ is a subalgebra of $H_{W(R)}(\nu)$.

Observe an important property of superalgebra $H_{W(R)}(\nu)$: the Lie superalgebra of its inner derivations[1] contains $\mathfrak{sl}_2$ generated by

$$T^{\alpha\beta} = \frac{1}{2} \sum_{i=1}^{N} \left\{ a_i^\alpha,\, a_i^\beta \right\}$$

which commute with $\mathbb{C}[W(R)]$, i.e., $[T^{\alpha\beta},\, R_{\vec{v}}] = 0$, and act on a_i^α as on $\mathfrak{sl}_2$-vectors:

$$\left[T^{\alpha\beta},\, a_i^\gamma \right] = \varepsilon^{\alpha\gamma} a_i^\beta + \varepsilon^{\beta\gamma} a_i^\alpha. \tag{4}$$

The restriction of operator T^{01} in the representation (3) on the subspace of $W(R)$-invariant functions on V is a second-order differential operator which is the well-known Hamiltonian of the rational Calogero model [6] based on the root system R [7]. One of the relations (4), namely, $[T^{01},\, a_i^\alpha] = -(-1)^\alpha a_i^\alpha$, allows one to find the wave functions of the equation $T^{01}\psi = \epsilon\psi$ via the usual Fock procedure with the vacuum $|0\rangle$ such that $a_i^0|0\rangle = 0$ for any i [5]. After $W(R)$-symmetrization these wave functions become the wave functions of Calogero Hamiltonian.

2 Supertraces on $H_{W(R)}(\nu)$

Definition 3. *Any linear complex-valued function* $str(\cdot)$ *on the superalgebra* $\mathcal{A}$ *such that*

$$str(fg) \;=\; (-1)^{\pi(f)\pi(g)} str(gf)$$

for any $f, g \in \mathcal{A}$ *with definite parity* $\pi(f)$ *and* $\pi(g)$ *is called a supertrace.*

---

[1] Let $\mathcal{A}$ be arbitrary associative superalgebra. Then, the operators $\mathcal{D}_x$ which act on $\mathcal{A}$ via $\mathcal{D}_x(y) = [x,\, y\}$ (supercommutator) constitute the Lie superalgebra of *inner* derivations.

The restriction of every supertrace on $\mathbb{C}[W(R)]$ is completely determined by its values on $W(R) \subset \mathbb{C}[W(R)]$ and the function *str* is a central function on $W(R)$, i.e., it is constant on the conjugacy classes.

Let us introduce the grading E on the vector space of $\mathbb{C}[W(R)]$. Consider the subspace $\mathcal{E}(g) \subset V$:

$$\mathcal{E}(g) = \{x \in V \mid \quad gx = -xg\} \text{ for } g \in W(R).$$

Set

$$E(g) = dim\,\mathcal{E}(g).$$

The following theorem was proved in [8]

Theorem 1. *The number of linearly independent supertraces $Q(R)$ on $H_{W(R)}(\nu)$ does not depend on ν and is equal to the number of conjugacy classes in $W(R)$ with $E(g) = 0$.*

This Theorem helps to find the number $Q(R)$ for an arbitrary root system R.

Theorem 1 implies, evidently, the following statement

Theorem 2.

$$Q(R_1 \oplus R_2) = Q(R_1)Q(R_2).$$

Therefore, the problem of finding $Q(R)$ is reduced to the problem of finding $Q(R)$ for irreducible root systems R.

3 The numbers $Q(R)$ for irreducible root systems

R	$Q(R)$	for the proof, see
A_{n-1}	the number of partitions of n into a sum of odd positive integers	[9]
B_n, C_n, BC_n	the number of partitions of n into a sum of positive integers	[10]
D_n	the number of partitions of n into a sum of positive integers with even number of even integers	[10]
E_6	9	Appendix 1
E_7	12	Appendix 2
E_8	30	Appendix 3
F_4	9	Appendix 4
G_2	3	[11]
H_3	4	Appendix 5
H_4	20	Appendix 6
$I_2(n)$	$\left[\frac{n+1}{2}\right]$	[11]

Appendices

Appendix 1. E_6

The conjugacy classes of the Weyl group E_6 are described in Table 9 of [12]. The following 9 classes have not the root -1:

$$\phi,\ A_2,\ A_2^2,\ A_4,\ D_4(a_1),\ A_2^3,\ E_6,\ E_6(a_1),\ E_6(a_2).$$

Appendix 2. E_7

The conjugacy classes of the Weyl group E_7 are described in Table 10 of [12]. The following 12 classes have not the root -1:

$$\phi,\ A_2,\ A_2^2,\ A_4,\ D_4(a_1),\ A_2^3,\ A_4\times A_2,\ A_6,\ D_6(a_1),\ E_6,\ E_6(a_1),\ E_6(a_2).$$

Appendix 3. E_8

The conjugacy classes of the Weyl group E_8 are described in Table 11 of [12]. The following 30 classes have not the root -1:

$$\begin{array}{l}\phi,\ A_2,\ A_2^2,\ A_4,\ D_4(a_1),\ A_2^3,\ A_4\times A_2,\ A_6,\ D_4(a_1)\times A_2,\ D_6(a_1),\\ E_6,\ E_6(a_1),\ E_6(a_2),\ A_2^4,\ A_4^2,\ A_8,\ D_4(a_1)^2,\ D_8(a_1),\ D_8(a_3),\\ E_6\times A_2,\ E_6(a_2)\times A_2,\ E_8,\ E_8(a_i)\ (i=1,\ldots 8).\end{array}$$

Appendix 4. F_4

The conjugacy classes of the Weyl group F_4 are described in Table 8 of [12]. The following 9 classes have not the root -1:

$$\phi,\ A_2,\ \tilde{A}_2,\ B_2,\ A_2\times\tilde{A}_2,\ D_4(a_1),\ B_4,\ F_4,\ F_4(a_1).$$

Appendix 5. H_3

Let $k=(\sqrt{5}+1)/2$. Then the reflections

$$\begin{aligned} a &= \begin{pmatrix} 1 & 0 & 0\\ 0 & -1 & 0\\ 0 & 0 & 1\end{pmatrix}\\ b &= \frac{1}{2}\begin{pmatrix} 1 & k & k-1\\ k & 1-k & -1\\ k-1 & -1 & k\end{pmatrix}\\ c &= \begin{pmatrix} -1 & 0 & 0\\ 0 & 1 & 0\\ 0 & 0 & 1\end{pmatrix}\end{aligned}$$

corresponding to the roots $\vec{e}_2$, $\frac{1}{2}(-\vec{e}_1 + k\vec{e}_2 + k^{-1}\vec{e}_3)$ and $\vec{e}_1$ satisfy the relations

$$a^2 = b^2 = c^2 = 1, \quad (ab)^5 = (bc)^3 = (ac)^2 = 1$$

and define the Coxeter group H_3.

As $H_3 = S_5^e \times C_2$ (See [1], Ch.VI, Sect. 4, Ex.11 d), p. 284; [13], p.160 and references there), where S_5^e is the group of even permutations of 5 elements, $C_2 = \{1, -1\}$, ($|S_5^e| = 60$, $|H_3| = 120$), H_3 has 10 conjugacy classes, 5 with positive determinant and 5 with negative one.

The conjugacy classes with positive determinant are described by their representatives

The representative	The characteristic polynomial
1	$(1-t)^3$
ac	$(1-t)(1+t)^2$
bc	$(1-t)(t^2+t+1)$
ab	$(1-t)[t^2+(1-k)t+1]$
$abab$	$(1-t)(t^2+kt+1)$

Each of this characteristic polynomials has the root $+1$, and so each characteristic polynomials of any conjugacy class with negative determinant has the root -1. The conjugacy class with representative ac also has the root -1.

So, the number of conjugacy classes without root -1 is equal to 4.

Appendix 6. H_4

According to [13], all 34 conjugacy classes of H_4 are described by their representatives acting on the space of quaternions

$$g_{lr}: \quad x \mapsto lxr^* \tag{A6.1}$$

and

$$g_p^*: \quad x \mapsto px^*. \tag{A6.2}$$

and all 25 pairs of unit quaternions l and r and 9 unit quaternions p are listed in Table 3 of [13].

Each operator (A6.2) has the root -1. Indeed, the equation $px^* = -x$ has nonzero solution $x = -1 + p$ if $p \neq 1$, and x is an arbitrary imaginary quaternion if $p = 1$.

Each operator (A6.1) has not the root -1 if and only if

$$l_0 + r_0 \neq 0. \tag{A6.3}$$

Indeed, the determinant of the map $x \mapsto lx + xr$ is equal to $4(l_0 + r_0)^2$.

There are 20 pairs of l, r in Table 3 of [13] satisfying the condition (A6.3), namely, K_i with $i = 1$, 4, 6, 8, $10 - 25$.

References

[1] N.Bourbaki, ”Groupes et algèbres de Lie”, ch.IV-ch.VI, Hermann, Paris, 1968.

[2] James E.Humphreys, ”Reflection groups and Coxeter groups”, Cambridge University Press, 1990.

[3] C.F.Dunkl, Trans. Am. Math. Soc. **311** (1989) 167.

[4] A. Polychronakos, Phys. Rev. Lett. **69** (1992) 703.

[5] L. Brink, H. Hansson and M.A. Vasiliev, Phys. Lett. **B286** (1992) 109.

[6] F. Calogero, J. Math. Phys., **10** (1969) 2191, 2197; *ibid* **12** (1971) 419.

[7] M. A.Olshanetsky and A. M. Perelomov, Phys. Rep., **94** (1983) 313.

[8] S.E. Konstein, Supertraces on the Superalgebra of Observables of Rational Calogero Model based on the Root System, arXiv:hep-th/9803213.

[9] S.E. Konstein and M.A. Vasiliev, J. Math. Phys. **37** (1996) 2872.

[10] S.E. Konstein, Teor. Mat. Fiz., **111** (1997) 252-262.

[11] S.E. Konstein, Rational Calogero models based on rank-2 root systems: supertraces on the superalgebras of observables, arXiv:math/9801001.

[12] R.W.Carter, Conjugacy classes in the weyl group, Compositio Mathematica, **25** n^{o}1, (1972) 1-59.

[13] Larry C.Grove, ”The characters of the hecatonicosahedroidal group”, Journal für die reine und angewandte Mathematik, **265** (1974) 160-169.